

MASSIVE NEUTRINOS AND LEPTON-FLAVOUR-VIOLATING PROCESSES

J.A. CASAS

*Instituto de Estructura de la Materia, CSIC
Serrano 123, 28006 Madrid*

A. IBARRA

*Department of Physics, Theoretical Physics, University of Oxford
1 Keble Road, Oxford OX1 3NP, United Kingdom*

If neutrino masses and mixings are suitable to explain the atmospheric and solar neutrino fluxes, this amounts to contributions to FCNC processes, in particular $\mu \rightarrow e, \gamma$. If the theory is supersymmetric and the origin of the masses is a see-saw mechanism, we show that the prediction for $\text{BR}(\mu \rightarrow e, \gamma)$ is in general larger than the experimental upper bound, especially if the solar data are explained by a large angle MSW effect, which recent analyses suggest as the preferred scenario.

1 See-saw, RG-induced LFV soft terms and $l_i \rightarrow l_j, \gamma$

In the pure Standard Model, flavour is exactly conserved in the leptonic sector since one can always choose a basis in which the (charged) lepton Yukawa matrix, $\mathbf{Y}_e$, and gauge interactions are flavour-diagonal. If neutrinos are massive and mixed, as suggested by the observation of atmospheric and solar fluxes¹, this is no longer true and there exists a source of lepton flavour violation (LFV), in analogy with the Kobayashi–Maskawa mechanism in the quark sector. Unfortunately, due to the smallness of the neutrinos masses, the predicted branching ratios for these processes are so tiny that they are completely unobservable, namely $\text{BR}(\mu \rightarrow e\gamma) < 10^{-50}$ ².

In a supersymmetric (SUSY) framework the situation is completely different. Besides the previous mechanism, supersymmetry provides new direct sources of flavour violation in the leptonic sector, namely the possible presence of off-diagonal soft terms³. In a self-explanatory notation, they have the form

$$-\mathcal{L}_{\text{soft}} = (m_L^2)_{ij} \bar{L}_i L_j + (m_{e_R}^2)_{ij} \bar{e}_{Ri} e_{Rj} + (A_{eij} e_{Ri}^c H_1 L_j + \text{h.c.}) + \text{etc.}, \quad (1)$$

where we have written explicitly just the soft breaking terms in the leptonic sector, namely scalar masses and trilinear scalar terms. All the fields in the previous equation denote just the corresponding scalar components. Concerning flavour violation the most conservative starting point for $\mathcal{L}_{\text{soft}}$ is the assumption of universality, which corresponds to take

$$(m_L^2)_{ij} = m_0^2 \mathbf{1}, \quad (m_{e_R}^2)_{ij} = m_0^2 \mathbf{1}, \quad A_{eij} = A_0 \mathbf{Y}_{eij}, \quad (2)$$

so that working in the L_i and e_{Ri} basis where $\mathbf{Y}_e$ is diagonal, the soft terms do not contain off-diagonal (lepton flavour violating) entries.

It turns out, however, that even under this extremely conservative assumption, if neutrinos are massive, radiative corrections may generate off-diagonal soft terms.

The most interesting example of this occurs when neutrino masses are produced by a (supersymmetric) see-saw mechanism⁴. This is based upon a superpotential

$$W = W_0 - \frac{1}{2} \nu_R^c{}^T \mathcal{M} \nu_R^c + \nu_R^c{}^T \mathbf{Y}_\nu L \cdot H_2, \quad (3)$$

where W_0 is the observable superpotential, except for neutrino masses, of the preferred version of the supersymmetric SM, e.g. the MSSM. The extra terms involve three additional neutrino chiral fields (one per generation; indices are suppressed) not charged under the SM group: ν_{Ri} ($i = e, \mu, \tau$). $\mathbf{Y}_\nu$ is the matrix of neutrino Yukawa couplings, L_i ($i = e, \mu, \tau$) are the left-handed lepton doublets and H_2 is the hypercharge +1/2 Higgs doublet. The Dirac mass matrix is given by $\mathbf{m}_D = \mathbf{Y}_\nu \langle H_2^0 \rangle$. Finally, $\mathcal{M}$ is a 3×3 Majorana mass matrix whose natural scale, say M , is much larger than the electroweak scale or any soft mass. Below M the theory is governed by an effective superpotential $W_{eff} = W_0 + \frac{1}{2} (\mathbf{Y}_\nu L \cdot H_2)^T \mathcal{M}^{-1} (\mathbf{Y}_\nu L \cdot H_2)$, obtained by integrating out the heavy neutrino fields in (3). Hence, the effective neutrino mass matrix, $\mathcal{M}_\nu$, is given by

$$\kappa \equiv \mathcal{M}_\nu / \langle H_2^0 \rangle^2 = \mathbf{Y}_\nu^T \mathcal{M}^{-1} \mathbf{Y}_\nu, \quad (4)$$

where $\langle H_2^0 \rangle^2 = v_2^2 = v^2 \sin^2 \beta$ and $v = 174$ GeV. The experimental data about neutrino masses and mixings are referred to the $\mathcal{M}_\nu$ matrix, or equivalently κ , evaluated at low energy (electroweak scale)^a.

Turning back to the structure of the SUSY soft-breaking terms, the universality condition (2) can only be imposed at a certain scale, typically at the scale at which the soft breaking terms are generated, e.g. M_X in GUT models. Below that scale, the RGEs of the soft terms, which contain non-diagonal contributions proportional to $\mathbf{Y}_\nu^+ \mathbf{Y}_\nu$, induce off-diagonal soft terms^{5,6,7,8}. These contributions are decoupled at the characteristic scale of the right-handed neutrinos, M . More precisely, in the leading-log approximation^b, the off-diagonal soft terms at low-energy are given by

$$\begin{aligned} (m_L^2)_{ij} &\simeq \frac{-1}{8\pi^2} (3m_0^2 + A_0^2) (\mathbf{Y}_\nu^+ \mathbf{Y}_\nu)_{ij} \log \frac{M_X}{M}, \\ (m_{eR}^2)_{ij} &\simeq 0, \\ (A_e)_{ij} &\simeq \frac{-3}{8\pi^2} A_0 Y_i (\mathbf{Y}_\nu^+ \mathbf{Y}_\nu)_{ij} \log \frac{M_X}{M}, \end{aligned} \quad (5)$$

^aIt should be noted that eq.(4) is defined at the “Majorana scale”, M . Therefore, in order to compare to the experiment one has still to run κ down to low energy through the corresponding RGE.

^bWe use the leading-log approximation through the text in order to make the results easily understandable. Nevertheless, the numerical results, to be exposed below, have been obtained by integrating the full set of RGEs.




Figure 1. Dominant Feynman diagrams contributing to $l_i \rightarrow l_j, \gamma$ in the mass-insertion approximation. $\tilde{L}_i$ are the slepton doublets in the basis where the gauge interactions and the charged-lepton Yukawa couplings are flavour-diagonal. $\tilde{\chi}_A$ denote the charginos and neutralinos.

where $i \neq j$ and Y_{l_i} is the Yukawa coupling of the charged lepton l_i .

The previous off-diagonal soft terms induce LFV processes, like $l_i \rightarrow l_j, \gamma$. The precise form of $\text{BR}(l_i \rightarrow l_j, \gamma)$ that we have used in our computations is a rather cumbersome expression⁷. However, for the sake of the physical discussion it is interesting to think in the mass-insertion approximation to identify the dominant contributions. As discussed in ref.⁸, these correspond to the mass-insertion diagrams enhanced by $\tan \beta$ factors. All of them are proportional to $m_{L_{ij}}^2$, and have the generic form shown in Fig. 1. Thus the size of the branching ratios is given by

$$\begin{aligned} \text{BR}(l_i \rightarrow l_j, \gamma) &\sim \frac{\alpha^3}{G_F^2} \frac{|m_{L_{ij}}^2|^2}{m_S^8} \tan^2 \beta \\ &\sim \frac{\alpha^3}{G_F^2 m_S^8} \left| \frac{-1}{8\pi^2} (3m_0^2 + A_0^2) \log \frac{M_X}{M} \right|^2 |(\mathbf{Y}_\nu^+ \mathbf{Y}_\nu)_{ij}|^2 \tan^2 \beta \quad (6) \end{aligned}$$

where we have used eqs.(5). The $\mathbf{Y}_\nu^+ \mathbf{Y}_\nu$ matrix is therefore the crucial quantity for the computation of $\text{BR}(l_i \rightarrow l_j, \gamma)$. Hence, in order to make predictions on $\text{BR}(l_i \rightarrow l_j, \gamma)$ we need to determine the most general form of $\mathbf{Y}_\nu$ and $\mathbf{Y}_\nu^+ \mathbf{Y}_\nu$, compatible with all the phenomenological requirements. Recall that the latter are referred to the $\mathcal{M}_\nu$ matrix, evaluated at low energy, rather than to $\mathbf{Y}_\nu$ itself. So, this is a non-trivial task that we discuss in the next section. Notice also the strong dependence of $\text{BR}(l_i \rightarrow l_j, \gamma)$ on $\tan \beta$ and the fact that the larger (smaller) the initial scale at which universality is imposed, the larger $\text{BR}(l_i \rightarrow l_j, \gamma)$.

2 General textures reproducing experimental data

Working in the flavour basis in which the charged-lepton Yukawa matrix, $\mathbf{Y}_e$, and gauge interactions are flavour-diagonal, the neutrino mass matrix, $\mathcal{M}_\nu$,

or equivalently the κ matrix defined in eq.(4), is diagonalized by the MNS ⁹ matrix U according to

$$U^T \kappa U = \text{diag}(\kappa_1, \kappa_2, \kappa_3) \equiv D_\kappa, \quad (7)$$

where U is a unitary matrix that relates flavour to mass eigenstates. It is possible, and sometimes convenient, to choose $\kappa_i \geq 0$. Then, U can be written as $U = V \cdot \text{diag}(e^{-i\phi/2}, e^{-i\phi'/2}, 1)$, where ϕ and ϕ' are CP violating phases (if different from 0 or π) and V has the ordinary form of a CKM matrix

$$V = \begin{pmatrix} c_{13}c_{12} & c_{13}s_{12} & s_{13}e^{-i\delta} \\ -c_{23}s_{12} - s_{23}s_{13}c_{12}e^{i\delta} & c_{23}c_{12} - s_{23}s_{13}s_{12}e^{i\delta} & s_{23}c_{13} \\ s_{23}s_{12} - c_{23}s_{13}c_{12}e^{i\delta} & -s_{23}c_{12} - c_{23}s_{13}s_{12}e^{i\delta} & c_{23}c_{13} \end{pmatrix}. \quad (8)$$

The experimental information about neutrinos consists of information about the low-energy spectrum of neutrinos, contained in D_κ , and about the neutrino mixing angles (and CP phases), contained in U . Let us discuss them in order.

The experimental (solar and atmospheric) data ¹ strongly suggest a hierarchy of neutrino mass-splittings, $\Delta\kappa_{sol}^2 \ll \Delta\kappa_{atm}^2$. Numerically, $\Delta\kappa_{atm}^2 \sim 3 \times 10^{-3} \text{eV}^2/v_2^4$ ¹⁰, while the value of $\Delta\kappa_{sol}^2$ depends on the solution considered to explain the solar neutrino problem ^{11,12}, i.e. large-angle, small angle or LOW MSW solutions (LAMSW, SAMSW and LOW respectively), or vacuum oscillations solution (VO). They require, in eV^2/v_2^4 units, $\Delta\kappa_{sol}^2 \sim 3 \times 10^{-5}$, 10^{-7} , 5×10^{-6} and 8×10^{-10} respectively. The most favoured one from the recent analyses of data ¹⁰ is the LAMSW. In any case, there are basically three types of neutrino spectra consistent with the hierarchy of mass-splittings ¹³: *hierarchical* ($\kappa_1^2 \ll \kappa_2^2 \ll \kappa_3^2$), *“intermediate”* ($\kappa_1^2 \sim \kappa_2^2 \gg \kappa_3^2$) and *“degenerate”* ($\kappa_1^2 \sim \kappa_2^2 \sim \kappa_3^2$). In the usual notation, $\Delta\kappa_{atm}^2 \equiv \Delta\kappa_{32}^2$, $\Delta\kappa_{sol}^2 \equiv \Delta\kappa_{21}^2$.

Concerning the mixing angles, θ_{23} and θ_{13} are constrained by the atmospheric and CHOOZ data to be near maximal and minimal, respectively. The θ_{12} angle depends on the solution considered for the solar neutrino problem: it should be either near maximal (LAMSW, LOW and VO) or near minimal (SAMSW). Hence, the two basic forms that U can present are either a single-maximal or (more plausibly) a bimaximal mixing matrix. Schematically,

$$U \sim \begin{pmatrix} 1 & 0 & 0 \\ 0 & \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ 0 & -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{pmatrix} \quad \text{or} \quad U \sim \begin{pmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} & 0 \\ -\frac{1}{2} & \frac{1}{2} & \frac{1}{\sqrt{2}} \\ \frac{1}{2} & -\frac{1}{2} & \frac{1}{\sqrt{2}} \end{pmatrix}, \quad (9)$$

Let us turn to our question of what is the most general form of $\mathbf{Y}_\nu$ and $\mathbf{Y}_\nu^+ \mathbf{Y}_\nu$ compatible with all the previous phenomenological requirements in a see-saw scenario. Notice, in the first place, that one can always choose to work in a basis of right neutrinos where $\mathcal{M}$ is diagonal

$$\mathcal{M} = \text{diag}(\mathcal{M}_1, \mathcal{M}_2, \mathcal{M}_3) \equiv D_{\mathcal{M}}, \quad (10)$$

with $\mathcal{M}_i \geq 0$. Then, from eqs.(4, 7), $D_\kappa = U^T \mathbf{Y}_\nu^T D_{\sqrt{\mathcal{M}-1}} D_{\sqrt{\mathcal{M}-1}} \mathbf{Y}_\nu U$, where, in an obvious notation, $D_{\sqrt{A}} \equiv +\sqrt{D_A}$. Consequently, the most general form of $\mathbf{Y}_\nu$ and $\mathbf{Y}_\nu^+ \mathbf{Y}_\nu$ is

$$\mathbf{Y}_\nu = D_{\sqrt{\mathcal{M}}} R D_{\sqrt{\kappa}} U^+ \quad (11)$$

$$\mathbf{Y}_\nu^+ \mathbf{Y}_\nu = U D_{\sqrt{\kappa}} R^+ D_{\mathcal{M}} R D_{\sqrt{\kappa}} U^+ \quad (12)$$

where R is any orthogonal matrix (R can be complex provided $R^T R = \mathbf{1}$).

So, besides the physical and measurable low-energy parameters, contained in D_κ and U , $\mathbf{Y}_\nu$ and $\mathbf{Y}_\nu^+ \mathbf{Y}_\nu$ depend on the three (unknown) positive mass eigenvalues of the righthanded neutrinos, contained in $D_{\mathcal{M}}$, and on the three (unknown) complex parameters defining R . We will see, however, that in practical cases the number of relevant free parameters becomes drastically reduced. Notice also that $\mathbf{Y}_\nu^+ \mathbf{Y}_\nu$, and therefore eq.(12), does not depend on the ν_R -basis, and thus on the fact that $\mathcal{M}$ is diagonal or not.

Two (very) special cases of eqs.(11, 12) occur when $R = \mathbf{1}$ and when $\mathbf{Y}_\nu$ has the form $\mathbf{Y}_\nu = W D_Y$, where D_Y is a diagonal matrix and W is a unitary matrix. Then, there exists a basis of L_i , ν_{Ri} where all the leptonic flavour violation arises from the $\mathbf{Y}_e$ (i.e. from the charged leptons) or from the $\mathcal{M}$ (i.e. the right neutrinos) matrices respectively. In the second case $\mathbf{Y}_\nu^+ \mathbf{Y}_\nu$ is diagonal, so the predictions for $\text{BR}(l_i \rightarrow l_j, \gamma)$ are negligible.

Next, we study the general predictions for $\text{BR}(l_i \rightarrow l_j, \gamma)$, focussing on $\text{BR}(\mu \rightarrow e \gamma)$, by considering, in a separate way, some interesting scenarios that often appear in the literature. A more detailed discussion can be found in Ref.¹⁴.

3 Predictions for $\text{BR}(l_i \rightarrow l_j, \gamma)$

ν_L 's and ν_R 's completely hierarchical

In this case $D_\kappa \simeq \text{diag}(0, \kappa_2, \kappa_3)$, $D_{\mathcal{M}} \simeq \text{diag}(0, 0, \mathcal{M}_3)$.

★ If R is a generic matrix, with $R_{32} \neq 0$ or $R_{33} \neq 0$, $(\mathbf{Y}_\nu^+ \mathbf{Y}_\nu)_{ij}$ is given by

$$(\mathbf{Y}_\nu^+ \mathbf{Y}_\nu)_{ij} \sim (\mathbf{Y}_\nu)_{3i}^* (\mathbf{Y}_\nu)_{3j} \sim \mathcal{M}_3 \left[\sum_{l=2,3} R_{3l}^* \sqrt{\kappa_l} U_{il} \right] \left[\sum_{l'=2,3} R_{3l} \sqrt{\kappa_{l'}} U_{jl'}^* \right]$$

Parameterizing R as

$$R = \begin{pmatrix} \hat{c}_2 \hat{c}_3 & -\hat{c}_1 \hat{s}_3 - \hat{s}_1 \hat{s}_2 \hat{c}_3 & \hat{s}_1 \hat{s}_3 - \hat{c}_1 \hat{s}_2 \hat{c}_3 \\ \hat{c}_2 \hat{s}_3 & \hat{c}_1 \hat{c}_3 - \hat{s}_1 \hat{s}_2 \hat{s}_3 & -\hat{s}_1 \hat{c}_3 - \hat{c}_1 \hat{s}_2 \hat{s}_3 \\ \hat{s}_2 & \hat{s}_1 \hat{c}_2 & \hat{c}_1 \hat{c}_2 \end{pmatrix}, \quad (13)$$




Figure 2. $\text{BR}(\mu \rightarrow e, \gamma)$ vs. the unknown angle $\hat{\theta}_1$ for the case of hierarchical (left and right) neutrinos, a typical set of supersymmetric parameters, and different values of the largest neutrino Yukawa coupling, Y_0 , at the “unification” scale, M_X . Y_t denotes the value of the top Yukawa coupling at that scale. θ_1 is taken real for simplicity, so the two limits $\hat{\theta}_1 = 0, \pi$ of the horizontal axis represent the same physical point. The dashed lines correspond to the present and forthcoming experimental upper bounds¹⁵.

where $\hat{\theta}_1, \hat{\theta}_2, \hat{\theta}_3$ are arbitrary complex angles [eq.(13) is sufficiently general for this case], one obtains in particular

$$(\mathbf{Y}_\nu^\dagger \mathbf{Y}_\nu)_{21} \sim \frac{|Y_0|^2}{|\hat{s}_1|^2 \kappa_2 + |\hat{c}_1|^2 \kappa_3} \hat{c}_1^* \hat{s}_1 \sqrt{\kappa_3 \kappa_2} U_{23} U_{12}^* \quad (14)$$

Here $|Y_0|^2$ is the largest eigenvalue of $\mathbf{Y}_\nu^\dagger \mathbf{Y}_\nu$ and $\hat{\theta}_1$ is an arbitrary complex angle. The branching ratio just depends on $|Y_0|^2$ and $\hat{\theta}_1$.

For the LAMSW scenario the previous equation generically gives $\text{BR}(\mu \rightarrow e, \gamma)$ above the *present* experimental limits, at least for $Y_0 = \mathcal{O}(1)$, as it occurs in the unified scenarios. (This holds until $m_0 > 1.5$ TeV, except for a narrow range at low m_0 .) Actually, even for $Y_0 = \mathcal{O}(10^{-1})$, most of the parameter space will be probed in the forthcoming generation of experiments experiments. This is illustrated in Fig. 2.

★ The only exceptions to the previous result are

- If $\hat{\theta}_1$ is such that $(\mathbf{Y}_\nu)_{31} \simeq 0$ or $(\mathbf{Y}_\nu)_{32} \simeq 0$. Then $(\mathbf{Y}_\nu^\dagger \mathbf{Y}_\nu)_{21} \simeq 0$ and $\text{BR}(\mu \rightarrow e, \gamma)$ is small. These two special values of $\hat{\theta}_1$ are visible in Fig. 2 and correspond to $\tan \hat{\theta}_1 \simeq -\sqrt{\frac{\kappa_3}{\kappa_2}} \frac{V_{13}^*}{V_{12}^*} \simeq 0$ and $\tan \hat{\theta}_1 \simeq -\sqrt{\frac{\kappa_3}{\kappa_2}} \frac{V_{23}^*}{V_{22}^*}$. The associated textures in our basis are

$$\mathbf{Y}_\nu \propto \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & V_{31} & -V_{21} \end{pmatrix} \propto \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 1 & 1 \end{pmatrix}, \quad (15)$$

$$\mathbf{Y}_\nu \propto \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ -V_{31} & 0 & V_{11} \end{pmatrix} \propto \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ -1 & 0 & \sqrt{2} \end{pmatrix} . \quad (16)$$

where the numerical values of the entries correspond to a bimaximal mixing V matrix. For these cases $(\mathbf{Y}_\nu^\dagger \mathbf{Y}_\nu)_{21} \propto \mathcal{M}_2$ (instead $\propto \mathcal{M}_3$), which can still be sizeable. In any case other processes, as $\text{BR}(\tau \rightarrow \mu, \gamma)$, are not suppressed, normally lying above the forthcoming experimental upper bound.

- If R is such that $\mathbf{Y}_\nu^\dagger \mathbf{Y}_\nu$ is diagonal. This requires a very special form of R , which in particular has $R_{32}, R_{33} \simeq 0$.

ν_L 's hierarchical and ν_R 's degenerate

In this case $D_\kappa \simeq \text{diag}(0, \kappa_2, \kappa_3)$, $D_\mathcal{M} \simeq \text{diag}(\mathcal{M}, \mathcal{M}, \mathcal{M})$.

- ★ If R is real, this scenario is very predictive. Then $(\mathbf{Y}_\nu^\dagger \mathbf{Y}_\nu)_{ij}$ does not depend on R

$$(\mathbf{Y}_\nu^\dagger \mathbf{Y}_\nu)_{ij} = \mathcal{M} \kappa_l U_{il} U_{lj}^\dagger \simeq \mathcal{M} [\kappa_2 U_{i2} U_{j2}^* + \kappa_3 U_{i3} U_{j3}^*] , \quad (17)$$

In particular, taking into account $U_{13} \simeq 0$,

$$(\mathbf{Y}_\nu^\dagger \mathbf{Y}_\nu)_{21} \simeq \mathcal{M} \kappa_2 U_{22} U_{12}^* = |Y_0|^2 \frac{\kappa_2}{\kappa_3} U_{22} U_{12}^* , \quad (18)$$

where $|Y_0|^2 \simeq \mathcal{M} \kappa_3$ is the largest eigenvalue of $\mathbf{Y}_\nu^\dagger \mathbf{Y}_\nu$.

The corresponding $\text{BR}(\mu \rightarrow e, \gamma)$ is illustrated in Fig. 3 for the LAMSW scenario.

The branching ratio turns out to be already above the present experimental limits except for a rather small region of m_0 values which should be probed by the next generation of experiments. Here are not special textures where the branching ratio becomes suppressed.

- ★ If R is complex, the analysis is more involved since it contains more arbitrary parameters. But in general the conclusion is the same: $\text{BR}(\mu \rightarrow e, \gamma)$ is at least of the same order as in the real case. Now there exists, however, the possibility of a (fine-tuned) cancellation.

ν_L 's quasi-degenerate

In this case $D_\kappa \simeq \text{diag}(\kappa_1, \kappa_2, \kappa_3)$, with $\kappa_1 \sim \kappa_2 \sim \kappa_3 \equiv \kappa$. Then, it is logical to assume that $\mathcal{M}$ has degenerate eigenvalues, otherwise a big conspiracy would be needed between $\mathbf{Y}_\nu$ and $\mathcal{M}$. Hence $D_\mathcal{M} \simeq \text{diag}(\mathcal{M}, \mathcal{M}, \mathcal{M})$.

- ★ If R is real, $\mathbf{Y}_\nu^\dagger \mathbf{Y}_\nu = \mathcal{M} U D_\kappa U^\dagger$. In particular, since $U_{13} \simeq 0$,

$$(\mathbf{Y}_\nu^\dagger \mathbf{Y}_\nu)_{21} = \mathcal{M} [U_{21} U_{11}^* (\kappa_1 - \kappa_2)] = |Y_0|^2 U_{21} U_{11}^* \frac{\Delta \kappa_{sol}^2}{\kappa^2} . \quad (19)$$




Figure 3. $\text{BR}(\mu \rightarrow e, \gamma)$ vs. the universal scalar mass, m_0 , for the case of hierarchical (degenerate) left (right) neutrinos, R real and typical sets of supersymmetric parameters. The dashed lines correspond to the present and forthcoming upper bounds. A top-neutrino “unification” condition has been used to fix the value of the largest neutrino Yukawa coupling at high energy. The curves do not fall below the present bound until $m_0 > 1.6$ TeV.

where, again, $|Y_0|^2 \simeq \mathcal{M}\kappa$ is the largest eigenvalue of $\mathbf{Y}_\nu^\dagger \mathbf{Y}_\nu$. This equation is identical to eq.(18), multiplied by $\kappa^{-2} \sqrt{\Delta\kappa_{sol}^2 \Delta\kappa_{atm}^2}$. This is a factor $\sim 10^{-4}$ for the LAMSW. Therefore all the plots representing $\text{BR}(\mu \rightarrow e, \gamma)$ in the previous scenario (Fig. 3) are valid here, but with the vertical axis re-scaled eight orders of magnitude smaller. Consequently, $\text{BR}(\mu \rightarrow e, \gamma)$ is naturally suppressed below the present (and even forthcoming) limits.

- ★ If R is complex, $\mathbf{Y}_\nu^\dagger \mathbf{Y}_\nu = \mathcal{M} U D_{\sqrt{\kappa}} R^\dagger R D_{\sqrt{\kappa}} U^\dagger$, which may have sizeable off-diagonal entries. Hence, $\text{BR}(\mu \rightarrow e, \gamma)$, could be very large in this case.
- ★ If the (quasi-) degeneracy is only partial: $\kappa_3 \ll \kappa_1 \simeq \kappa_2 \equiv \kappa \sim \sqrt{\Delta\kappa_{atm}^2}$, $(\mathbf{Y}_\nu^\dagger \mathbf{Y}_\nu)_{21}$ is given (for R real) by eq.(18), multiplied now by $\sqrt{\frac{\Delta\kappa_{sol}^2}{\Delta\kappa_{atm}^2}}$. This represents a suppression factor $\sim 10^{-1}$ for the LAMSW, which means that Fig. 3 should be re-scaled by a factor $\sim 10^{-2}$. As a consequence, $\text{BR}(\mu \rightarrow e, \gamma)$ for this partially degenerate scenario should be testable within the next generation of experiments. The conclusion is similar for $\text{BR}(\tau \rightarrow \mu, \gamma)$.

For generic complex R , the value of $\text{BR}(\mu \rightarrow e, \gamma)$ does not get any suppression and falls naturally above the present experimental limits.

4 Conclusions

If the origin of the neutrino masses is a supersymmetric see-saw, which is probably the most attractive scenario to explain their smallness, then the leptonic soft breaking terms acquire off-diagonal contributions through the RG running, which drive non-vanishing $\text{BR}(l_i \rightarrow l_j, \gamma)$. These contributions are proportional to $(\mathbf{Y}_\nu^\dagger \mathbf{Y}_\nu)_{ij}$, where $\mathbf{Y}_\nu$ is the neutrino Yukawa matrix,

Therefore, in order to make predictions for these branching ratios, one has first to determine the most general form of $\mathbf{Y}_\nu$ and $\mathbf{Y}_\nu^\dagger \mathbf{Y}_\nu$, compatible with all the phenomenological requirements. This is summarized in eqs.(11, 12).

Then, we have shown that the predictions for $\text{BR}(\mu \rightarrow e, \gamma)$ are normally *above* the *present* experimental limits if the three following conditions occur

1. The solution to the solar neutrino problem is the LAMSW, as favoured by the most recent analyses.
2. $Y_0(M_X) = \mathcal{O}(1)$, where $|Y_0|^2$ is the largest eigenvalue of $(\mathbf{Y}_\nu^\dagger \mathbf{Y}_\nu)$. This occurs e.g. in most grand-unified scenarios.
3. The soft-breaking terms are generated at a high-energy scale, e.g. M_X , above the Majorana mass of the right-handed neutrinos, M .

These conditions are very plausible. In our opinion, the most natural scenarios fulfill them, but certainly there exists other possibilities. E.g. it may happen that supersymmetry is broken at a scale below M . This is the case of gauge-mediated scenarios, where there would be no generation of off-diagonal leptonic soft terms through the RG running.

Even under the previous 1–3 conditions, there are physical scenarios compatible with the present $\text{BR}(\mu \rightarrow e, \gamma)$ experimental limits. Namely

- Whenever all the leptonic flavour violation can be attributed to the sector of right-handed neutrinos. In this case there is no RG generation of non-diagonal soft terms.
- In the scenario of hierarchical (left and right) neutrino masses, *if* $\mathbf{Y}_\nu$ has (in our basis) one of the two special textures shown in eqs.(15, 16).
- If the left-handed neutrinos are quasi-degenerate and the R matrix in eq.(11) is real.

In our opinion, the scenario of quasi-degenerate neutrinos and the one with gauge mediated supersymmetry breaking represent the most plausible explanations to the absence of $\mu \rightarrow e, \gamma$ observations, specially if the absence persists after the next generation of experiments.

As a final conclusion, the discovery of neutrino oscillations makes much more plausible the possibility of observing lepton-flavour-violation processes, specially $\mu \rightarrow e, \gamma$, if the theory is supersymmetric and the neutrino masses are generated by a see-saw mechanism. Large regions of the parameter space are *already excluded* on these grounds, and there exists great chances to observe $\mu \rightarrow e, \gamma$ in the near future (PSI, 2003). This means that, hopefully, we will have signals of supersymmetry before LHC.

References

1. See e.g. talks by Y. Oyama and P. Harris in this conference.
2. S. M. Bilenkii, S. T. Petcov and B. Pontecorvo, *Phys. Lett. B* **67** (1977) 309; T. P. Cheng and L. Li, *Phys. Rev. Lett.* **45** (1980) 1908; W. J. Marciano and A. I. Sanda, *Phys. Lett. B* **67** (1977) 303; B. W. Lee, S. Pakvasa, R. E. Shrock and H. Sugawara, *Phys. Rev. Lett.* **38** (1977) 937 [Erratum-ibid. **38** (1977) 937].
3. See e.g. talks by A. Masiero and D. Carvalho in this conference.
4. M. Gell-Mann, P. Ramond and R. Slansky, proceedings of the Supergravity Stony Brook Workshop, New York, 1979, eds. P. Van Nieuwenhuizen and D. Freedman (North-Holland, Amsterdam); T. Yanagida, proceedings of the Workshop on Unified Theories and Baryon Number in the Universe, Tsukuba, Japan 1979 (edited by A. Sawada and A. Sugamoto, KEK Report No. 79-18, Tsukuba); R. Mohapatra and G. Senjanović, *Phys. Rev. Lett.* **44** (1980) 912, *Phys. Rev. D* **23** (1981) 165.
5. F. Borzumati and A. Masiero, *Phys. Rev. Lett.* **57** (1986) 961.
6. See e.g. J. Ellis, M. E. Gomez, G. K. Leontaris, S. Lola and D. V. Nanopoulos, data,” *Eur. Phys. J. C* **14** (2000) 319; M. E. Gomez, G. K. Leontaris, S. Lola and J. D. Vergados, *Phys. Rev. D* **59** (1999) 116009; D. F. Carvalho, M. E. Gomez and S. Khalil, hep-ph/0101250; R. Barbieri, L. Hall and A. Strumia, *Nucl. Phys. B* **445** (1995) 219.
7. J. Hisano, T. Moroi, K. Tobe and M. Yamaguchi, *Phys. Rev. D* **53** (1996) 2442.
8. J. Hisano and D. Nomura, *Phys. Rev. D* **59** (1999) 116005.
9. Z. Maki, M. Nakagawa and S. Sakata, *Prog. Theor. Phys.* **28** (1962) 870.
10. See e.g. M. C. Gonzalez-Garcia, M. Maltoni, C. Pena-Garay and J. W. Valle, *Phys. Rev. D* **63** (2001) 033005.
11. B. Pontecorvo, *Sov. Phys. JETP* **26** (1968) 984.
12. S. P. Mikheev and A. Y. Smirnov, *Sov. J. Nucl. Phys.* **42** (1985) 913; L. Wolfenstein, *Phys. Rev. D* **17** (1978) 2369.
13. G. Altarelli and F. Feruglio, *Phys. Lett. B* **439** (1998) 112, *JHEP* **9811** (1998) 021, and *Phys. Lett. B* **451** (1999) 388.
14. J. A. Casas and A. Ibarra, hep-ph/0103065.
15. M. L. Brooks *et al.* [MEGA Collaboration], *Phys. Rev. Lett.* **83** (1999) 1521; L. M. Barkov *et al.*, Research Proposal for an experiment at PSI

(1999), see the webpage: <http://meg.psi.ch/>.